

Quintron® Instructions

COST INFORMATION:

- The Quintron test is ~\$210 (through Snoqualmie Valley Hospital or Genova) if you do not have insurance or your insurance does not cover the cost of the test.
- Here is a link to a videos and instructions for the breath testing <https://www.doctorlelaaltman.com/resources>

SOME INSTRUCTIONS IN THE VIDEOS DIFFER FROM THESE INSTRUCTIONS, PLEASE DEFAULT TO THESE INSTURCTIONS

TEST PREPARATION¹:

1. For 1-2 weeks prior to the test – **no antibiotics, runny diarrhea (unless this is normal for you) or abnormal diarrhea pattern or barium studies**
2. No **colonoscopy for 4 weeks** prior to testing
3. For 4-5 days prior to the test – **no probiotics or laxatives**
4. For 1-day prior to test – please **stop any unnecessary supplements** the day before the test and resume after you complete the test. Essential medications can be continued.
5. For 12-24 hours* prior to the test – **please eat only the following foods:** (If you are diabetic, vegetarian or vegan, please see below.)
 - **White** rice (plain steamed)-only if this is something you tolerate well
 - **Eggs**
 - **Baked** or broiled chicken, fish or turkey
 - **Clear meat broth** (no bone broth or vegetable broth)
 - **Salt and pepper** only
 - **Water, Black tea and sBlack coffee** (without any creamer)
6. For 12 hours prior to the test – **fasting stage**, water ONLY during this stage *If you have fewer than 2 bowel movements per week or a redundant colon, then you will need to do the prep diet for **two days**.

OTHER PREP OPTIONS IF NECESSARY¹:

- - Diabetic If difficulty with fasting overnight, eat 1/4 of an unripe (green) banana or 2-3 baby carrots.
- - Vegetarian – eggs, white rice/wheat, hard aged cheese (small amount) and same diabetic approved foods as well.
- - Vegan – fasting for 24 hours is possible if blood sugar regulation is not an issue, additionally, can have white rice/bread, diabetics approved foods in addition to up to 2/3 cup plain/firm tofu (not silken)

MORNING OF THE TEST:

- - Must be awake for 1 hour prior to starting the test.

- For the morning of the test and during the test: no smoking, no sleeping, no eating, and no exercising

COMPLETING THE TEST:

Step 1: Fill out the labels for the collection tubes, making sure to fill out the **sample number, date and time**. Follow the instructions that come with your kit on collection of breath into bags. Be sure to exhale a normal breath (not a deep breath) when you are doing the breath collection samples.

Step 2: You can choose to either use mouthwash before collecting every sample or not at all. PLEASE DO NOT USE MOUTHWASH JUST ONCE AT THE BEGINNING OF THE TEST, THIS WILL ALTER YOUR RESULTS. The benefit of using mouthwash is that it may reduce the impact mouth bacteria play in the test results. If you have gingivitis, it is best to use mouthwash. If using mouthwash, please rinse your mouth prior to each sample. Move the mouthwash around the mouth for 30 seconds, forcing it between teeth and gargling before spitting it out. Take care not to swallow any mouthwash. Do a second rinse with water afterwards. Repeat this step just before each subsequent breath sample.

Step 3: Mix the in 8-10 oz of water for lactulose test - **DO NOT DRINK IT YET**, set it aside. You will need to pick this up from your pharmacy ahead of time.

Step 4: Collect first (baseline) breath specimen in collection tube (staple your collection bag according to your weight if you weigh 100 lbs or less, load the collection tube into the barrel with the gray rubber stick (there is a needle inside this) without puncturing the seal on the collection tube, blow through the mouthpiece to gently inflate the bag, and while you are still exhaling, push the collection tube through so that the seal is punctured by the needle. Remove the tube after 1-2 seconds.)

Step 5: Drink the preparation you just made immediately.

Step 6: Collect subsequent breath samples on this schedule: Sample #2: 20 minutes after baseline sample

Sample #3: 40 minutes after baseline sample

Sample #4: 60 minutes after baseline sample

Sample #5: 75 minutes after baseline sample Sample #6: 90 minutes after baseline sample

Sample #7: 110 minutes after baseline sample Sample #8: 130 minutes after baseline sample

Sample #9: 150 minutes after baseline sample Sample #10: 180 minutes after baseline sample

Step 7: Package up test and return to company